

Comment on: Dynamic and Static Properties of the Randomly Pinned Flux Array

Heiko Rieger

Institut für Theoretische Physik, Universität zu Köln, 50926 Köln, Germany

HLRZ c/o Forschungszentrum Jülich, Postfach 1913, 52425 Jülich, Germany

(February 13, 1995)

We reconsider the model of the randomly pinned planar flux array (or the two-dimensional XY-model with random fields and no vortices) in extensive Monte Carlo simulations. We show that contrary to what has been reported recently [Phys. Rev. Lett. **72**, 4133 (1994)] the disorder-averaged displacement correlation function is different from the pure case for larger values of the disorder. Its slope as a function of the logarithm of disorder is compatible with newer RG-calculation results.

In a recent letter Batrouni and Hwa [1] reported results of numerical simulations of the planar flux array described by the random phase model

$$H = \frac{\kappa}{2} \sum_{\langle ij \rangle} (u_i - u_j)^2 - \lambda \sum_i \cos(2\pi u_i - \beta_i) , \quad (1)$$

where $\langle ij \rangle$ denote nearest neighbor sites on a square lattice, u_i a real-valued displacement-like field, β_i a random phase uniformly distributed in the interval $[0, 2\pi]$, and λ the strength of the pinning potential. The main conclusion of the investigation in [1] was **a)** that the various renormalization group predictions existing up to that time could be ruled out by their numerical results, which I agree with, and **b)** that the disorder averaged correlation function $C(r) = [\langle (u_i - u_{i+\mathbf{r}})^2 \rangle]_{\text{av}}$, where $\langle \dots \rangle$ means the thermodynamic expectation value and $[\dots]_{\text{av}}$ the disorder average, is indistinguishable from the pure case, i.e. $C(r) = C_{\lambda=0}(r) = T/\kappa\pi \log r$ for $T \leq T_g = \kappa/\pi$.

In this comment I would like to point out that the last statement is a consequence of the weakness of the disorder they used, namely $\lambda = 0.15$, and that for stronger disorder (or larger length scales) the correlation function $C(r)$ differs significantly from the pure case. Furthermore the numerical data I obtain are compatible with the analytic predictions of [2,3] and the recent numerical results of a related model [4] with infinite disorder.

I used the usual Monte Carlo algorithm to calculate the static expectation values of the model (1). By setting $\kappa = 2$ the glass transition temperature is $T_g = 2/\pi \approx 0.637$. The system sizes were $L=32$ and 64 (where 1280 and 256 samples were used, respectively) and periodic boundary conditions were imposed, which forces the correlation function in the x -(and y)-direction to be symmetric $C(r) = C(L - r)$, $r \leq L/2$. By applying a least square fit of the data points for $C(r)$ with $r \leq L/4$ (where the effect of the periodic boundary conditions is still negligible) to the function $C(r) = a + b(T) \log r$ (T being the temperature) one gets the results shown in figure 1 for different disorder strengths.

For weak disorder $\lambda \sim 0.15$ the slope $b(T)$ is indeed indistinguishable from the pure case $b(T) = T/\kappa\pi$, as Batrouni and Hwa observed [1]. However, by increasing the disorder ($\lambda \geq 0.5$) one obtains a slope $b(T)$ that is significantly different from the pure case already

in the vicinity of T_g . Since systems with larger disorder are hard to equilibrate, only data for $\lambda \leq 2$ are shown, but the trend seems to be obvious: the estimate of $b(T)$ obtained from intermediate length scales increases with increasing disorder strength.

One should point out that $b(T)$ shown in figure 1 can only be interpreted as the slope of $C(r)$ as a function of $\log r$ for small length scales. For certain values of T and λ one observes a significant upward bend in these curves already at length scale that smaller than $L/4$, indicating that the asymptotic behavior is not yet reached. In figure 2 the derivative of $C(r)$ with respect to $\log r$, thus yielding a local slope $b(T, r)$, is depicted for $\lambda = 0.5$ and $T = 0.4$. This plot gives further support for the hypothesis that $\lim_{r \rightarrow \infty} b(T, r) = T_g/2\pi$, however, a quadratic dependency $C(r) \sim \log^2 r$, implying $\lim_{r \rightarrow \infty} b(T, r) = \infty$, cannot be excluded. With this observation in mind the data shown in figure 1 have to be interpreted as *lower bounds* for the asymptotic slope $b(T)$ of $C(r)$ as function of $\log r$. Hence the results for the correlation function are compatible with $b(T) = T_g/\kappa\pi$ for $T \leq T_g$, however, also a quadratic dependency $C(r) \sim \log^2 r$, implying $\lim_{r \rightarrow \infty} b(T, r) = \infty$, although unlikely, cannot be strictly excluded.

Concluding I have presented numerical evidence that the disorder averaged spatial correlation function of model (1) is indeed distinct from the pure case $\lambda = 0$. For weak disorder this becomes manifest only on length scales that are not attainable via Monte Carlo simulations yet, which is the reason why Batrouni and Hwa [1] were not able to detect it. However, my results agree with their conclusion that the various RG predictions, prior to their work, were incorrect.

I would like to thank T. Nattermann, S. Scheidl, L. H. Tang and J. Kierfeld for many valuable suggestions and discussions. The computations were done on the Parsytec-GCel1024 from the Center of Parallel Computing (ZPR) in Köln and the Intel Paragon System from the Supercomputercenter (HLRZ) at the Forschungszentrum Jülich. This work was performed within the SFB 341 Köln–Aachen–Jülich.

REFERENCES

- [1] G. G. Batrouni and T. Hwa, Phys. Rev. Lett. **72**, 4133 (1994).
- [2] S. E. Korshunov, Phys. Rev. B **48**, 3969 (1993).
- [3] T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **71**, 1530 (1994); preprint cond-mat/9501087 (1995).
- [4] D. Cule and Y. Shapir, Phys. Rev. Lett. **74**, 114 (1995); E. Marinari, R. Monasson and J. Ruiz-Lorenzo, preprint cond-mat/9503074.

FIGURES




FIG. 1. The slope $b(T)$, obtained via a least square fit of the MC-data for $r \leq L/4$, as a function of the temperature T . From top to bottom one has $\lambda = 2.0$ ($\bullet$), 1.0 ($\circ$), 0.5 ($\diamond$) and 0.25 ($\square$). Data for $\lambda = 0.15$, as obtained by Batrouni and Hwa [1] fall onto the thick full line, which is the analytic result for the pure model: $b(T) = T/\kappa\pi$. The dash-dotted line indicates the glass transition temperature $T_g = \kappa/\pi$ and the dotted line is at $b(T) = T_g/\kappa\pi$.




FIG. 2. The local slope $b(T, r) = [C(r + 1) - C(r)] / [\log(r + 1) - \log(r)]$ as a function of distance. Shown are data for three different equilibration times: $t_w = 3 \cdot 10^4$ ($\square$), $t_w = 10^5$ ($+$) and $t_w = 3 \cdot 10^5$ ($\diamond$). Curves for larger equilibration times are identical to the last one meaning that equilibration has been achieved. The dotted line is at $b = T_g / (\pi\kappa) = 1/\pi^2$. Because of the periodic boundary conditions of the finite system of length L one has $b(T, L/2) = 0$. If the prediction $C(\mathbf{r}) = T_g / (\pi\kappa) \log |\mathbf{r}|$ for $T \leq T_g$ holds, $b(T, r)$ has to approach this line for $r \rightarrow \infty$ in the infinite system.